

# Damvita-M

(Syrup – F. C. Tablets)

**COMPOSITION:**

Composition	Each 5 ml Syrup Contains:	Each F.Ctd. Tablet Contains:
Vit. A	3000 IU	5000 IU
Vit. D3	600 IU	400 IU
Vit. B1 (HCl)	3 mg	20 mg
Vit. B2 (phosphate)	6 mg	—
Vit. B2	—	10 mg
Vit. B6 (HCl)	2.5 mg	2 mg
Vit. B12	—	6 mcg
Vit. PP	10 mg	40 mg
Vit. C	75 mg	60 mg
Vit. E (acetate)	5 mg	50 mg
Folic Acid	—	3 mg
Ca – Panthenate	5 mg	20 mg
Iron (ferrous gluconate)	10.8 mg	—
Iron (ferrous fumarate)	—	18 mg
Magnesium (sulfate)	300 mcg	5 mg
Calcium (gluconate)	60 mg	—
Calcium (Phosphate)	—	100 mg
Copper (Sulfate)	300 mcg	500 mcg
Zinc (Sulfate)	0.5 mg	500 mcg

**PROPERTIES:**

**Damvita-M** is a multivitamin preparation with minerals. These vitamins and minerals are essential for normal metabolic functions including hematopoiesis. The B complex vitamins are necessary for the metabolism of carbohydrate, protein and fat. Ascorbic acid is involved in collagen formation and tissue restoration. Vitamin A is very necessary for normal retina function and for skin growth. Vitamin E is considered as antioxidant to keep normal function of Cell. Vitamins which are soluble in water (B, C vitamins) do not store at sufficient quantity in body, and excrete in large amount in urine so it should be compensated with **Damvita-M** to keep normal concentration in tissues. **Damvita-M** contains suitable concentration of vitamin B6 to correct the disorder of tryptophen metabolism to its normal value, this happens following estrogen taking as contraceptive pills. **Damvita-M** contains folic acid which is so important during pregnancy. Iron, Zinc, Copper are elements for enzyme structure to complete basic function of cells. Calcium is very important to normal growth of bones, teeth and for neuron cell function. Copper is important to prevent hypopigmentation of hair, Zinc is necessary for hair growth and protection of dry skin and night blindness (supplement agent to Vit. A), also helps the remedy of tissues quickly. Iron is necessary to the treatment and prevention of anemia.

**INDICATIONS:**

This medication is used for the prevention and treatment of multivitamin deficiency and the anemia caused by bad nutrition and unbalanced diet, the need for this medication increases greatly in these cases:

- In case of pregnancy and lactation.
- In case of using antibiotics against acute infections.
- In case of diarrhea for long time.
- In case of disorder of the process of carbohydrate protein and fat metabolism.
- In case of alcohol addiction.
- After surgery to help the remedy of tissues quickly.
- In case of abnormal growth.
- In case of tissues injuries like burns, bruises and wounds.
- In case of old age.

**CONTRAINdications:**

Patients allergic to the medication or for one of its components.

**DRUG INTERACTIONS:**

Patients who use levodopa.

**WARNING:**

This drug should be used cautiously in these cases:

- Hyper-calcemia.
- Increase of vitamins A and D.
- Patients with acute kidney failure or urolithiasis.
- Pregnant women should not exceed recommended doses, because of Vit. A.

**SIDE EFFECTS:**

It's rarely noticed when taken under physician's supervision, but some side effects may appear like: Allergic skin rashes, pruritus, vertigo and headache, feces of patient using iron may go black.

**PRECAUTIONS:**

- Keep medication out of reach of children.
- Store below 30°C.
- Use only under physician's supervision.

**DOSAGE:**

Syrup: Adults: (5–10) ml, 1–2 times daily.

Children: (2.5–5) ml, 1–2 times daily.

Coated Tablets: Adults: 1–3 coated tablets daily or as doctor prescribe.

**STORAGE:**

Store at room temperature (15–30)°C. Keep in dry place.

**PACKAGE:**

Carton box of 30 F.C. tablets blister packed.

Carton box of 125 ml glass bottle.

PK0045/01

TPP1301461	THIS IS A MEDICAMENT
– A medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.	
– Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold you the medicament.	
– The doctor and pharmacist are experts in medicine, its benefits and risks.	
– Do not by yourself interrupt the period of treatment prescribed for you.	
– Do not repeat the same prescription without consulting your doctor.	
– Keep medicaments out of reach of children.	

(Council of Arab Health Ministers & Arab Pharmacists Association)

**Ultra Medica** Pharmaceutical Industries: Sednaya - Damascus - Syria

Tel.: +963 11 5955339 – Fax: +963 11 5975174

E-mail: ultramed@scs-net.org – www.ultra-medica.net

